

BIBLIOGRAFIA (GENETICA Y MORFOLOGIA)

- ALCOVER, J.A., SANDERS, E. & SANCHIZ, B. (1984): El registro fósil de los sapos parteros (Anura, Discoglossidae) de Baleares. 109-121. En: HEMMER, H. & ALCOVER, J. A. (eds.). Història biològica del ferreret (Life history of the Mallorcan Midwife Toad). Ed. Moll. Ciutat de Mallorca.
- ALLENDORF, F.W. (1977): Electromorphs or alleles. Genetics 87: 821-822.
- ALLENDORF, F.W & PHELPS, S.R. (1981): Use of allelic frequencies to describe population structure. Can. J. Fish. Aquat. Sci. 38: 1507-1514.
- AQUADRO, C.F. & AVISE, J.C. (1982a): An assessment of "hidden" heterogeneity within electromorphs at three loci in deer mice. Genetics 102: 269-284.
- AQUADRO, C.F. & AVISE, J.C. (1982b): Evolutionary genetics of birds. VI. A reexamination of protein divergence using varied electrophoretic conditions. Evolution 36: 1003-1019.
- ARNTZEN, J.W. (1989 a): Biochemical systematics of midwife toads (genus *Alytes*): Comunicación oral. IV Congreso Nacional de Herpetología, Madrid.
- ARNTZEN, J.W. (1989 b): The identification of eggs of some species of Palearctic newts (genus *Triturus*) by starch gel electrophoresis. Isozyme Bull. 22: 69.
- ARNTZEN, J.W. (1990): Enzyme electrophoretic markers in the study of interspecific mate choice and individual mating success in newts. Isozyme Bull. 23: 79.
- ARNTZEN, J.W. & SZYMURA, J.M. (1984): Genetic differentiation between African and European midwife toads (*Alytes*, Discoglossidae). Bijdragen tot de Dierkunde 54(2): 157-162.
- AVISE, J.C. & AQUADRO, C.F. (1982): A comparative summary of genetic distances in the vertebrates. Evol. Biol. 15: 151-158.
- AVISE, J.C. (1974): Systematic value of electrophoretic data. Syst. Zool. 23: 465-481.
- AVISE, J.C. (1980): Diferenciación genética durante la especiación. 110-127. En: AYALA, F.J. (ed.). Evolución Molecular. Omega, Barcelona.
- AYALA, F.J. (1983): Genetic polymorphisms: from electrophoresis to DNA sequences. Experientia 39: 813-823.
- AYALA, F.J. & POWELL, J.R. (1972): Allozymes as diagnostic characters of sibling species of *Drosophila*. Proc. Natl. Acad. Sci. 69: 1094-1096.
- AYALA, F.J.; POWELL, J.R.; TRACEY, M.L.; MOURAO, C.A. & PEREZ-SALAS, S. (1972): Enzyme variability in the *Drosophila willistoni* group. IV. Genic variation in natural populations of *Drosophila willistoni*. Genetics 70: 113-139.
- BARBADILLO, L.J. (1987): La guía de INCAFO de los anfibios y reptiles de la Península Ibérica, Islas Baleares y Canarias. Ed. Incafo, 694 págs.
- BEA, A. (1985): Atlas de los anfibios y reptiles de Alava, Vizcaya y Guipúzcoa. 15-24. En: ALVAREZ, J., BEA, A., FAUS, J. M., CASTIEN, E. & MENDIOLA, I. Atlas de los vertebrados continentales de Alava, Vizcaya y Guipúzcoa. Gobierno Vasco.

BOGART, J.P., LICHT, L.E., OLDHAM, M.J. & DARBYSHIRE, S.J. (1985): Electrophoretic identification of *Ambystoma laterale* and *Ambystoma texanum* as well as their diploid and triploid interspecific hybrids (Amphibia: Caudata) on Pelee Island, Ontario. Can. J. Zool. 63: 340-347.

BOSCA, E. (1879 a): Nueva especie de Alytes en España. Actas, An. Soc. Esp. Hist. Nat. 8: 22.

BOSCA, E. (1879 b): Alytes cisternasii, descripción de un nuevo batracio de la fauna española. An. Soc. Esp. Hist. Nat. 8: 217-227.

BREWER, G.J. (1970): An Introduction Isozyme Techniques. Academic Press, Nueva York.

BUSACK, S.D. (1977): Zoogeography of amphibians and reptiles in Cádiz province, Spain. Ann. Carnegie Mus. 46: 285-316.

BUSACK, S.D. (1986): Biochemical and morphological differentiation in Spanish and Moroccan population of *Discoglossus* and the description of a new species from Southern Spain (Amphibia, Anura, Discoglossidae). Ann. Carnegie Museum 55: 41-61.

BUTH, D.G. (1979): Genetic relationships among the torrent suckers, genus *Thoburnia*. Biochem. Syst. Ecol. 3: 311-316.

BUTH, D.G. (1984): The application of electrophoretic data in systematic studies. Annu. Rev. Ecol. Syst. 15: 501-522.

CAMIN, J.H. & SOKAL, R.R. (1965): A method for deducing branching sequences in phylogeny. Evolution 19: 311-326.

CAPULA, M., NASCETTI, G., LANZA, B., BULLINI, L. & CRESPO, E.G. (1985): Morphological and genetic differentiation between the Iberian and the other west Mediterranean *Discoglossus* species (Amphibia Salientia Discoglossidae). Monitore zool. ital. (N.S.) 19: 69-90.

CASE, S.M. (1978 a): Biochemical systematics of members of the genus *Rana* native to western North America. Syst. Zool. 27: 299-311.

CASE, S.M. (1978 b): Electrophoretic variation in two species of ranid frogs, *Rana boylei* and *R. muscosa*. Copeia 1978: 311-320.

CASE, S.M., HANELINE, P.G. & SMITH, M. F. (1975): Protein variation in several species of *Hyla*. Syst. Zool. 24: 281-295.

CAVALLI-SFORZA, L.L. & EDWARDS, A.W.F. (1967): Phylogenetic analysis: Models and estimation procedures. Evolution 21: 550-570.

CEI, J.M. (1980): Amphibians of Argentina. Monitore zool. ital. (N.S.) Monogr. 2: 1-609.

CLARKE, B.T. (1984): General skeletal morphology. 45-59. En: HEMMER, H. & ALCOVER, J. A. (eds). Història biològica del ferreret (Life history of the Mallorcan Midwife Toad). Ed. Moll. Ciutat de Mallorca.

CORBETT, K. (1989): Conservation of European Reptiles & Amphibians. Ed. Helm, Londres. 274 págs.

COYNE, J. (1982): Gel electrophoresis and cryptic protein variation, pp. 1-32. En: RATTazzi, M., SCANDALIOS, J. & WHITT, G. (eds). Isozymes: Current Topics in Biological and Medical Research, vol. 6. Molecular Structure and Regulation. Liss, Nueva York.

CRACRAFT, J. (1974): Phylogeny and evolution of the ratite birds. Ibis 116: 494-521.

CRESPO, E.G. (1976): Contribuição para o estudo das espécies ibéricas do género Alytes (Alytes

cisternasii Boscá e Alytes obstetricans boscai Lataste)-(Amphibia, Salientia): Bolm. Soc. Port. Cienc. Nat. 17: 39-54.

CRESPO, E.G. (1979): Contribuiç_o para o conhecimento da biologia dos Alytes ibéricos, Alytes obstetricans boscai Lataste, 1879 e Alytes cisternasii Boscá, 1879 (Amphibia, Salientia): A problemática da especiaç_o de Alytes cisternasii. Tese. Universidade de Lisboa, Faculdade de Ciéncias, Lisboa. 1,2: 399.

CRESPO, E.G. (1981 a): Contribuç_o para o conhecimento da la biología das espécies ibéricas de Alytes, Alytes obstetricans boscai Lataste, 1879 e Alytes cisternasii Boscá, 1879 (Amphibia, Discoglossidae): tegumento (histología e polipeptidos activos). Arquivos do Museo Bocage 1(2) Sér. C: 33-56.

CRESPO, E.G. (1981 b): Contribuç_o para o conhecimento da la biología das espécies ibéricas de Alytes, Alytes obstetricans boscai Lataste, 1879 e Alytes cisternasii Boscá, 1879 (Amphibia, Discoglossidae): regulaç_o hidrica-balanco osmótico. Arquivos do Museo Bocage 1(4) Sér. C: 77-132.

CRESPO, E.G. (1982): Contribuç_o para o conhecimento da la biología das espécies ibéricas de Alytes, Alytes obstetricans boscai Lataste, 1879 e Alytes cisternasii Boscá, 1879 (Amphibia, Discoglossidae): morfología dos adultos e dos girinos. Arquivos do Museo Bocage 1(7) Sér. C: 255-312.

CRESPO, E.G. (1985): Sur la structure génétique de deux "populations" allopatriques d'Alytes obstetricans boscai et d'Alytes cisternasii (Amphibia, Discoglossidae) du Portugal. Alytes 4(1): 1-11.

CRESPO, E.G.; VIEGAS, A.M. & VICENTE, L.A. (1984): Sur quelques caractéristiques biochimiques de Baleaphryne muletensis. 199-212. En: HEMMER, H. & ALCOVER, J. A. (eds.). Història biològica del ferreret (Life history of the Mallorcan Midwife Toad). Ed. Moll. Ciutat de Mallorca.

CROWLEY, Th. & NORTH, G.R. (1991): Paleoclimatology. Oxford University Press, Nueva York. 399 págs.

DUBOIS, A. (1983): Classification et nomenclature suprageneric des amphibiens anoures. Bull. Men. Soc. Linnéenne Lyon 52: 270-276.

DUBOIS, A. (1987): Again on the nomenclature of frogs. Alytes, 6(1-2): 27-55.

DUELLMAN, W.E. & TRUEB, L. (1985): Biology of Amphibians. Ed. McGraw-Hill Book Company. 670 págs.

ESTES, R. & SANCHIZ, B. (1982): New Discoglossid and Paleobatrachid frogs from the Late Cretaceous of Wyoming and Montana, and a Review of other frogs from the Lance and Hell Creek Formations. Journal of Vertebrate Paleontology 2(1): 9-20.

FARRIS, J.S. (1970): Methods for computing Wagner trees. Syst. Zool. 34: 21-34.

FARRIS, J.S. (1972): Estimating phylogenetic trees from distance matrices. Am. Natur. 106: 645-668.

FARRIS, J.S. (1977): Phylogenetic analysis under Dollo's Law. Syst. Zool. 26: 77-88.

FARRIS, J.S. (1980): The information content or the phylogenetic system. Syst. Zool. 28: 483-519.

FARRIS, J.S. (1981): Distance data in phylogenetic analysis, 2-23. En: FUNK, V.A. & BROOKS, D.R. (eds). Advances in Cladistics. New York Botanical Garden, Nueva York.

FELSENSTEIN, J. (1984): Distance methods for inferring phylogenies: A justification. Evolution 38: 16-24.

FELSENSTEIN, J. (1989): PHYLIP, phylogenetic inference package, version 3.2. Department of

Genetics, SK 50. University of Washington, Seattle, Washington, USA.

FIALA, K.L. & SOKAL, R.S. (1985): Factors determining the accuracy of cladogram estimation: Evaluation using computer simulation. *Evolution* 39: 609-622.

FITCH, W.M. & ATCHLEY, W.R. (1987): Divergence in inbred strains of mice: a comparison of three different types of data. 203-216. En: PATTERSON, C. (ed.). *Molecules and morphology in evolution: conflict or compromise?*. Cambridge University Press.

FITCH, W.M. & MARGOLIASH, E. (1967): Construction of phylogenetic trees. *Science* 155: 279-284.

FITCH, W.M. (1971): Toward defining the course of evolution: Minimal change for a specific tree topology. *Syst. Zool.* 20: 406-416.

FREIFELDER, D. (1982): *Physical Biochemistry: Applications to Biochemistry and Molecular Biology*. 2nd. ed. Freeman, Nueva York.

FRETEY, J. (1975): *Guide des Reptiles et Batraciens de France*. Ed. Hatier, París. 239 págs.

FROST, D.R. (ed.) (1985): *Amphibian species of the world : a taxonomic and geographical reference*. Allen Press & The Association of Systematics Collections. Lawrence.

GARCIA-PARIS, M., ASTUDILLO, G., PRIETO, J. & MARQUEZ, R. (1990): Distribución de *Alytes cisternasii* Bosca, 1879, en el centro de la Península Ibérica. *Rev. Esp. Herp.* 4: 87-91.

GILLESPIE, J.H. & KOJIMA, K. (1968): The degree of polymorphism in enzymes involved in energy production compared to that in monospecific enzymes in two *Drosophila ananassae* populations. *Genetics* 61: 582-585.

GOLLMANN, G. (1984): Allozymic and morphological variation in the hybrid zone between *Bombina bombina* and *Bombina variegata* (Anura, Discoglossidae) in northeastern Austria. *Z. Zool. Syst. Evolut.-forsch.* 22: 51-64.

GOOD, D.A., WURST, G.Z. & WAKE, D.B. (1987): Patterns of Geographic Variation in Allozymes of the Olympic Salamander, *Rhyacotriton olympicus* (Caudata: Dicamptodontidae): *Fieldiana: Zoology* N.S. 32: 1-15.

GOTTLIEB, L.D. (1982): Conservation and duplication of isozymes in plants. *Science* 216: 373-380.

GOULD, S.J. (1982): The meaning of punctuated equilibrium and its role in validating a hierarchical approach to macroevolution. En: MILKMAN, R. (ed.). *Perspectives in evolution*. Sinauer, Sunderland. USA.

GREEN, D.M. (1986): Systematics and evolution of western north american frogs allied to *Rana aurora* and *Rana boylii*: electrophoretic evidence. *Syst. Zool.* 35(3): 283-296.

HAMES, B. D. & RICKWOOD, D. (Eds) (1981): *Gel Electrophoresis of Proteins: A practical Approach*. IRL Press, Oxford.

HARLAND, W.B. et al. (1990): *A geological time scale 1989*. Cambridge University Press, Cambridge. 263 págs.

HARRIS, H. & HOPKINSON, D. A. (1976): *Handbook of Enzyme Electrophoresis in Human Genetics*. North-Holland, Amsterdam.

HARRIS, H. (1966): Enzyme polymorphism in man. *Proc. Roy. Soc. London, ser B.*, 164: 298-310.

HAYASI, T. & MATSUI, M. (1988): Biochemical differentiation in Japanese newts, genus *Cynops*

(Salamandridae). Zool. Sci. 5: 1121-1136.

HAYASI, T., MATSUI, M., UTSUNOMIYA, T., TANAKA, S. & OHTA, H. (1992): Allozyme variation in the newt *Tylototriton andersoni* from three islands of the Ryukyu Archipelago. Herpetologica 48(2): 178-184.

HEMMER, H. (1975): Serumeiweibunterschied zwischen *Bufo b. bufo* und *Bufo bufo spinosus* (Salientia, Bufonidae). Salamandra 11(1): 61-63.

HEMMER, H. (1984): The Mallorcan midwife toad -*Alytes (Baleaphryne) muletensis*-: a new phylogenetic level within a conservative genus. 235-243 págs. En: HEMMER, H. & ALCOVER, J. A. (eds). Història biològica del ferreret (Life history of the Mallorcan Midwife Toad). Ed. Moll. Ciutat de Mallorca.

HEMMER, H. & ALCOVER, J.A. (eds) (1984): Història biològica del ferreret. Moll. Ciutat de Mallorca.

HEMMER, H.; KADEL, B. & KADEL, K. (1981): The Balearic toad (*Bufo viridis balearicus* (Boettger, 1881)), human bronze age culture, and Mediterranean biogeography. Amph. Rept., 2: 217 - 230.

HENNING, W. (1950): Grundzüge einer Theorie der phylogenetischen Systematik. Deutscher Zentralverlag, Berlin.

HENNING, W. (1966): Phylogenetic systematics. University of Illinois Press, Urbana.

HERRERO, P. (1984): Estudios citogenéticos en *Baleaphryne muletensis*. 223-230. En: HEMMER, H. & ALCOVER, J. A. (eds). Història biològica del ferreret (Life history of the Mallorcan Midwife Toad). Moll. Ciutat de Mallorca.

HIGHTON, R. (1989): Biochemical evolution in the slimy salamanders of the *Plethodon glutinosus* complex in the eastern United States. Part. I. Geographic protein variation. Univ. Illinois Biol. Monogr. 57: 1-78.

HIGHTON, R. & WEBSTER, T.P. (1976): Geographic protein variation and divergence in populations of the salamander *Plethodon cinereus*. Evolution 30: 33-45.

HILLIS, D.M. (1984): Misuse and modification of Nei's genetic distance. Syst. Zool. 33: 238-240.

HILLIS, D.M. (1987): Molecular versus morphological approaches to systematics. Ann. Rev. Ecol. Syst. 18: 23-42.

HUBBY, J. L. & LEWONTIN, R.C. (1966): A molecular approach to the study of genic heterozygosity in natural populations. I. The number of alleles at different loci in *Drosophila pseudoobscura*. Genetics 54: 577- 594.

HUBBY, J. L. & THROCKMORTON, L.H. (1965): Protein differences in *Drosophila*. II. Comparative species genetics and evolutionary problems. Genetics 52: 203-215.

HUNTER, R.L. & MARKERT, C.L. (1957): Histochemical demonstration of enzymes separated by zone electrophoresis in starch gels. Science 125: 1294-1295.

JACOBS, J.F. (1987): A preliminary investigation of geographic genetic variation and systematics of the two-lined salamander, *Eurycea bislineata* (Green). Herpetologica 43(4): 423-446.

JOHNSON, G.B. (1977): Assessing electrophoretic similarity: The problem of hidden heterogeneity. Annu. Rev. Ecol. Syst. 8: 309-328.

KALEZIC, M.L. & TUCIC, N. (1984): Genic diversity and population genetic structure of *Triturus*

vulgaris (Urodela, Salamandridae): Evolution 38: 389-401.

KIMURA, M. (1983 a): The neutral theory of molecular evolution. En: NEI, M. & KOHEN, R. K. (eds): Evolution of Genes and Proteins. Sinauer, Sunderland. USA.

KIMURA, M. (1983 b): The neutral theory of molecular evolution. Cambridge Univ. Press., Cambridge.

KIMURA, M. & OHTA, T. (1971): Protein polymorphism as a phase of molecular evolution. Nature 229: 467-469.

KING, J.L. & OHTA, T. (1975): Polyallelic mutational equilibria. Genetics 79: 681-691.

KLUGE, A.G. (1983): Cladistics and the classification of the great ape. 151-177. En: CIOCHAN, R.L. & CORRUCINI, R.S. (eds). New Interpretations of Ape and Human Ancestry. Plenun, Nueva York.

KLUGE, A.G. & FARRIS, J.S. (1969): Quantitative phyletics and the evolution of anurans. Syst. Zool. 18: 1-32.

KOBAYASI, T., MILNER, G.B., TEEL, D. & UTTER, F.M. (1984): Genetic basis for electrophoretic variation of adenosine deaminase in chinook salmon. Trans. Am. Fish. Soc. 113: 86-89.

LANZA, B., CEI, J. M. & CRESPO, E. G. (1975): Immunological evidence for the specific status of *Discoglossus pictus* Otth, 1837 and *D. sardus* Tschudi, 1837, with notes on the families Discoglossidae Günther, 1858 and Bombinidae Fitzinger, 1826 (Amphibia Salientia). Monitore zool. ital. (N.S.) 9(3-4): 153-162.

LANZA, B., CEI, J.M. & CRESPO, E.G. (1976): Further immunological evidence for the validity of the family Bombinidae (Amphibia Salientia). Monitore zool. ital. (N.S.) 10(3): 311-314.

LANZA, B., NASCETTI, G. & BULLINI, L. (1982): Tassonomia biochimica del genere *Hydromantes* (Amphibia, Plethodontidae). Boll. Zool., Atti XLIX Convegno Unione Zoologica Italiana 49(suppl.): 103.

LANZA, B., NASCETTI, G., CAPULA, M. & BULLINI, L. (1984): Genetic relationships among west Mediterranean *Discoglossus* with the descriptions of a new species (Amphibia Salientia Discoglossidae): Monitore zool. ital. (N.S.) 18: 133-152.

LANZA, B., NASCETTI, G., CAPULA, M. & BULLINI, L. (1986): Les discoglosses de la région Méditerranéenne occidentale (Amphibia; Anura; Discoglossidae). Bull. Soc. Herp. Fr. 40: 16-27.

LARSON, A., WAKE, D.B. & YANEV, K.P. (1984): Measuring gene flow among populations having high levels of genetic fragmentation. Genetics 106: 293-308.

LATASTE, F. (1879): Sur une nouvelle forme de Batracien anoure d_Europe (*Alytes obstetricans boscai*, n. subsp.) Rev. Int. Sci. Paris 4: 543-545.

LE QUESNE, W.J. (1974): The uniquely evolved character concept and its cladistic application. Syst. Zool. 23: 513-517.

LEMEN, C. (1983): The effectiveness of methods of shape analysis. Fieldiana, Zoology (N.S. 15) 1343: 1-17.

LEWONTIN, R.C. (1974): The Genetic Basis of Evolutionary Change. Columbia Univ. Press, Nueva York.

LEWONTIN, R.C. & HUBBY (1966): A molecular approach to the study of genic heterozygosity in natural populations. II Amounts of variation and degree of heterozygosity in natural populations of *Drosophila pseudoobscura*. Genetics 54: 595-609.

LIBIS, B. (1985): Nouvelle donnée sur la répartition au Maroc du crapaud accoucheur *Alytes maurus* Pasteur et Bons 1962. (Amphibia, Discoglossidae). Bull. Soc. Herp. Fr. 33: 52-53.

MANCHENKO, G.P. (1988): Subunit structure of enzymes: Allozymic data. Isozyme Bull. 21: 144-158.

MARKERT, C.L. (ed.) (1975): Isozymes. I. Molecular Structure. II, Physiological function. III, Developmental Biology. IV, Genetic and Evolution. Academic Press, New York, San Francisco, Londres.

MARKERT, C.L. (1983): Isozymes: Conceptual history and biological significance, pp. 1-17. En: RATTAZZI, M.C., SCANDALIOS, J.G. & WHITT, G.S. (eds). Isozymes: Current Topics in Biological and Medical Research, vol. 7. Molecular Structure and Regulation. Liss, Nueva York.

MARQUEZ, R. (1990): The mating systems of the midwife toads *Alytes obstetricans* and *Alytes cisternasii*. Dissertation in candidacy for the degree of Doctor of Philosophy. University of Chicago, USA.

MAXSON, L.R. (1984): Relationships of Baleaphryne to the discoglossid genera *Alytes*, *Bombina*, and *Discoglossus*. 193-198. En: HEMMER, H. & ALCOVER, J. A. (eds). Història biològica del ferreret (Life history of the Mallorcan Midwife Toad). Moll. Ciutat de Mallorca.

MAXSON, L.R. & MAXSON, R.D. (1979): Comparative albumin and biochemical evolution in plethodontid salamander. Evolution 33(4): 1057-1068.

MAXSON, L.R. & SZYMURA, J.M. (1984): Relationships among discoglossid frog: An albumin perspective. Amphibia - Reptilia 5: 245-252.

MAYOL, J. & ALCOVER, J. A. (1981): Survival of Baleaphryne Sanchíz and Adrover 1979 (Amphibia: Anura: Discoglossidae) on Mallorca. Amphibia-Reptilia 1(3/4): 343-345.

MAYOL, J., ALCOVER, J.A., ALOMAR, G., POMAR, G. JURADO, J. & JAUME, D. (1980): Supervivència de Baleaphryne (Amphibia: Anura: Discoglossidae) a les muntanyes de Mallorca. Nota preliminar. Butll. Inst. Cat. Hist. Nat. 45 (Sec. Zool.) 3: 115-119.

MAYOL, J., ALCOVER, J.A., JAUME, D., ALOMAR, G., JURADO, J. & POMAR, G. (1984): Morfología externa dels adults de Baleaphryne muletensis. 15-20. En: HEMMER, H. & ALCOVER, J. A. (eds). Història biològica del ferreret (Life history of the Mallorcan Midwife Toad). Ed. Moll. Ciutat de Mallorca.

MCLELLAN, T. (1984): Molecular charge and electrophoretic mobility in cetacean myoglobins of known sequence. Biochem. Genet. 22: 181-200.

MELLADO, J. & MATEO, J.A. (1992): New records of Moroccan herpetofauna. Herpetological Journal 2: 58-61.

MERTENS, R. & MÜLLER, L. (1928): Liste der Amphibien und Reptilien Europas. Abhandlungen der Senckenbergischen Naturforschenden Gesellschaft 41(1): 1-62.

MERTENS, R. & WERMUTH, H. (1960): Die Amphibien und Reptilien Europas. Waldemar Kramer, Frankfurt am Main. 1-59.

MICKEVICH, M.F. & JOHNSON, M.S. (1976): Congruence between morphological and allozyme data in evolutionary inference and character evolution. Syst. Zool. 25: 260-270.

MOORE, D.W. & YATES, T.L. (1983): Rate of protein inactivation in selected animals following death. J. Wildl. Manag. 47: 1166-1169.

MORITZ, C & HILLIS, D.M. (1990): Molecular Systematics: Context and Controversies. 1-10. En:

HILLIS, D.M. & MORITZ, C. (eds). Molecular Systematics. Sinauer Associates, Inc., Massachusetts, USA.

MURPHY, R. W. & MATSON, R. H. (1986): Gene expression in the tuatara, *Sphenodon punctatus*. N. Z. J. Zool. 13: 573-581.

MURPHY, R. W., SITES, J. W., BUTH, J. R., BUTH, D. G. & HAUFLER, C. H. (1990): Proteins I: Isozyme electrophoresis. En: HILLIS, D. M. & MORITZ, C. (eds). Molecular Systematics. Sinauer Ass. Inc. USA.

NEI, M. (1972): Genetic distance between populations. Am. Natur. 106: 283-292.

NEI, M. (1978): Estimation of average heterozygosity and genetic distance from a small number of individuals. Genetics 89: 583-590.

NEI, M. (1987): Molecular Evolutionary Genetics. Columbia Univ. Press, Nueva York.

NEI, M. & CHESSER, R.K. (1983): Estimation of fixation indices and gene diversification. Ann. Human Genet. 47: 253-259.

NELSON, G.J. (1974): Classification as an expression of phylogenetic relationships. Syst. Zool. 22: 344-359.

NEVO, E., BEILES, A. & BEN-SHLOMO, R. (1984): The evolutionary significance of genetic diversity: Ecological, demographic and life history correlates. Lec. Notes Biomath. 53: 13-213.

NISHIOKA, M., SUMIDA, M., OHTA, S. & SUZUKI, H. (1987): Speciation of three allied genera, *Buergeria*, *Rhacophorus* and *Polypedates*, elucidated by the method of electrophoretic analyses. Sci. Rep. Lab. Amphib. Biol. Hiroshima Univ. 9: 53-96.

PASTEUR, A., PASTEUR, G., BONHOMME, F., CATALAN, J. & BRITTON-DAVIDIAN, J. (1987): Manuel technique de génétique par électrophorèse des protéines. Ed. Technique et Documentation (Lavoisier), París. 1-217.

PASTEUR, G. (1985): Les paramètres statistiques communément utilisés dans l'exploitation des résultats de l'électrophorèse des protéines et leur avenir en systématique. En: GOYFFON, M. & D'HONDT, J.L. (eds). Electrophorèse et Taxinomie. Mémoires de la Société Zoologique de France, París: 141-180.

PASTEUR, G. & BONS, J. (1962): Note préliminaire sur l'*Alytes (obstetricans) maurus*: gemellarité ou polytopisme? Remarques biogeographiques, génétiques et taxonomiques. Bull. Soc. Zool. France 87: 71-79.

PASTEUR, G. & PASTEUR, N. (1980): Les critères biochimiques et l'espèce animale. En: Les Problèmes de l' Espèce dans le Règne Animal. C. Bocquet; J. Génermont et M. Lamotte (eds.), Mémoires de la Société Zoologique de France 40: 99-150.

PATTERSON, C. (ed.) (1987): Molecules and Morphology in Evolution: Conflict or Compromise?. Cambridge Univ. Press, Cambridge.

PLEGUEZUELOS, J.M. & MORENO, M. (1990): Atlas herpetológico de la provincia de Jaén. Junta de Andalucía. Agencia de Medio Ambiente. 75 págs.

RAFINSKI, J. & ARNTZEN, J.W. (1987): Biochemical systematics of the Old World newts, genus *Triturus*: allozyme data. Herpetologica 43: 446-457.

RICHARDSON, B.J., BAVERSTOCK, P.R. & ADAMS, M. (1986): Allozyme Electrophoresis. A handbook for animal Systematics and Population Structure. Academic Press, Sydney.

RIDER, C.C. & TAYLOR, C.B. (1980): Isoenzymes. Chapman & Hall, Londres.

ROGERS, J.S. (1972): Measures of genetic similarity and genetic distance. Studies in Genet. VII. Univ. Texas Publ. 7213: 145-153.

ROGERS, J.S. (1984): Deriving phylogenetic trees from allele frequencies: A comparison of nine generic distances. Syst. Zool. 35: 297-310.

ROGERS, J.S. (1986): Deriving phylogenetic trees from allele frequencies. Syst. Zool. 33: 52-63.

ROSA, H.D., VIEGAS, A.M. & CRESPO, E.G. (1990): Genetic structure of portuguese populations of midwife toads, with special reference to an isolate of *Alytes obstetricans*. Portugaliae Zoologica 1(3): 15-25.

SAITOU, N. & NEI, M. (1987): The neighbor-joining method: A new method for reconstructing phylogenetic trees. Mol. Biol. Evol. 4: 406-425.

SALVADOR, A. (1985): Guía de campo de los anfibios y reptiles de la Península Ibérica, Islas Baleares y Canarias. edición del autor 212 págs.

SANCHIZ, B. (1984): Análisis filogenético de la tribu Alytini (Anura, Discoglossidae) mediante el estudio de su morfoestructura ósea. 61-108. En: HEMMER, H. & ALCOVER, J. A. (eds.). Història biològica del ferreret (Life history of the Mallorcan Midwife Toad). Moll. Ciutat de Mallorca.

SANCHIZ, B. (1986): La evolución de los discoglósidos: implicaciones metodológicas. Resúmenes del Primer Congreso Nacional de Herpetología. Benicassim, Castellón.

SANCHIZ, B. & ADROVER, R. (1979): Anfibios fósiles del Pleistoceno de Mallorca. Doñana, Acta Vertebrata 4: 5-25.

SANCHIZ, B. & ALCOVER, J.A. (1982): Un nou Discoglòssid (Amphibia: Anura) de l'Holecè de Menorca. Butll. Inst. Cat. Hist. Nat. 48 (Sec. Geol. 3): 99-105.

SANCHIZ, B. & SCHLEICH, H.H. (1986): Revisión taxonómica de *Alytes grandis* Brunner (Amphibia, Anura). Estudios geol. 42: 471-473.

SANCHIZ, F.B. & VALDECASAS, A.G. (1980): Criterios metodológicos y glosario español de términos utilizados en sistemática cladística. Bol. R. Soc. Española Hist. nat. (Biol.), 78(1-2): 223-244.

SARICH, V.M. (1977): Rates, sample size, and the neutrality hypothesis for electrophoresis in evolutionary studies. Nature 265: 24-28.

SBORDONI, V., COBOLLI SBORDONI, M., DE MATTHAEIS, E., ALLEGRECCI, G., CESARONI, D., CACCONE, A. & CARPANETO, G.M. (1982): Orologi molecolari e paleogeografia: congruenza tra stime geocronologiche e datazioni elettroforetiche della divergenza nelle specie del genere *Euproctus* (Caudata, Salamandridae). Bollettino di Zoologia 49, suppl. 170: 170.

SCHLEE, D. (1971): Die Rekonstruktion der Phylogenetse mit Hennig's Prinzip. Waldemar Kramer, Frankfurt am Main.

SHAW, C.R. (1965): Electrophoretic variation in enzymes. Science 149: 936-943.

SHAW, C.R. & PRASAD, R. (1970): Starch gel electrophoresis of enzymes - a compilation of recipes. Biochem. Genet. 4: 297-330.

SIBLEY, C.G. & AHLQUIST, J.E. (1987). Avian phylogeny reconstructed from comparisons of the genetic material, DNA. 95-121. En: PATTERSON, C. (ed.). Molecules and Morphology in Evolution: Conflict or Compromise?. Cambridge Univ. Press, Cambridge.

SICILIANO, M.J. & SHAW, C.R. (1976): Separation and visualization of enzymes on gels. 185-209. En: SMITH, I. (ed.). Chromatographic and Electrophoretic Techniques, vol. 2. Wm. Heineman Medical Books, Londres.

SIMON, C.M. (1979): Evolution of periodical cicadas: Phylogenetic inferences based upon allozyme data. *Syst. Zool.* 28: 22-39.

SMITHIES, O. (1955): Zone electrophoresis in starch gels: Group variations in the serum proteins of normal individuals. *Biochem. J.* 61: 629-641.

SNEATH, P.H.A. & SOKAL, R.R. (1973): Numerical Taxonomy. W.H. Freeman, San Francisco.

SOKAL, R.R. & ROHLF, F.J. (1981): Biometry. 2nd ed. W.H. Freeman & Co., San Francisco. 859 págs.

SOULE, M.E. (1980): Variación aloenzimática: sus determinantes en el espacio y en el tiempo. 60-79. En: AYALA, F.J. (ed.). Evolución Molecular. Omega, Barcelona.

SWOFFORD, D.L. (1981): On the utility of the distance Wagner procedure, pp. 25-43. En: FUNK, V.A. & BROOKS, D.R. (eds.). Advances in Cladistics. Proc. First Meeting Willi Henning Soc. New York Botanical Gardens, Bronx, Nueva York.

SWOFFORD, D.L. & BERLOCHER, S.H. (1987): Inferring evolutionary trees from gene frequency data under the principle of maximum parsimony. *Syst Zool.* 36: 293-325.

SWOFFORD, D.L. & OLSEN, G.J. (1990): Phylogeny Reconstruction. 411-501. En: HILLIS, D.M. & MORITZ, C. (eds). Molecular Systematics. Sinauer, Sunderland. USA.

SWOFFORD, D.L. & SELANDER, R.B. (1981): BIOSYS-1: A FORTRAN program for the comprehensive analysis of electrophoretic data in population genetics and systematics. *J. Hered.* 72: 281-283.

SZYMURA, J.M. (1976): Hybridization between Discoglossid toads *Bombina bombina* and *Bombina variegata* in southern Poland as revealed by the electrophoretic technique. *Z. zool. Syst. Evolut.-forsch.* 14: 227-236.

SZYMURA, J.M. & BARTON, N.H. (1986): Genetic analysis of a hybrid zone between the fire-bellied toads, *Bombina bombina* and *B. variegata*, near Cracow in southern Poland. *Evolution* 40(6): 1141-1159.

THORPE, J.P. (1982): The Molecular Clock Hypothesis: Biochemical Evolution, Genetic Differentiation and Systematics. *Ann. Rev. Ecol. Syst.* 13: 139-168.

UZZELL, T. & BERGER, L. (1975): Electrophoretic phenotypes of *Rana ridibunda*, *Rana lessonae* and their hybridogenetic associate, *Rana esculenta*. *Proc. Acad. Nac. Sci. Philadelphia* 127: 13-23.

VIEGAS, A.M. & CRESPO, E.G. (1985): Sur la structure génétique de deux "populations" allopatriques d' *Alytes obstetricans boscai* et d' *Alytes cisternasii* (Amphibia, Discoglossidae) du Portugal. *Alytes* (París) 4: 1-11.

WAKE, D.B. & YANEV, K.P. (1986): Geographic variation in allozymes in a "ring species," the plethodontid salamander *Ensatina escholtzii* of western North America. *Evolution* 40(4): 702-715.

WASSERSUG, R.J. (1976): A procedure for differential staining of Cartilage and Bone in whole formalin-fixed vertebrates. *Stain Technology* 51(2): 131-134.

WILEY, E.O. (1981): Phylogenetics: The Theory and Practice of Phylogenetic Systematics. Wiley Interscience, Nueva York. 439 págs.

WILEY, E.O., SIEGEL-CAUSEY, D., BROOKS, D.R. & FUNK, V.A. (1991): The Compleat Cladist. A Primer of Phylogenetic Procedures. Special Publication No. 19: 1-158. University of Kansas, USA.

WILKINSON, L. (1989): SYSTAT: The System for Statistics. Evanston, IL: SYSTAT, Inc. 638 págs.

WILSON, A.C., CARLSON, S.S. & WHITE, T.J. (1977): Biochemical evolution. Annu. Rev. Biochem. 46: 573-639.

WILSON, A.C., SARICH, V.M. & MAXSON, L.R. (1974): The importance of gene rearrangement in evolution: Evidence from studies of rates of chromosomal, protein, and anatomical evolution. Proc. Natl. Aca. Sci. U.S.A. 71: 3028-3030.

WRIGHT, C.A. (ed.) (1978): Biochemical and Immunological Taxonomy of Animals. Academic Press, Londres.

ZAR, J.H. (1984): Biostatistical Analysis. (Second Edition). Prentice-Hall, New Jersey. 718 págs.