

VII. BIBLIOGRAFÍA

- ÁLVAREZ CÁCERES, R. (1995). *Estadística multivariable y no paramétrica con SPSS*. Díaz de Santos S.A.
- BALNEAVES, J.M. and DE LA MARE, P.J. (1989). *Root patterns of Pinus radiata on five ripping treatments in a canterbury forest*. N. Z. J. For. Sci. 19 (1): 29-40.
- BEYELER, J. (1996). *Effect of planting methods on field performance of Black Spruce five years after planting*. Forest Research Report 62: 1-16.
- BISQUERRA ALZINA, R. (1989). *Introducción conceptual al análisis multivariable* (Vol.1). PPU. Barcelona
- BRISSTTE J.C.; BARNETT J.P. (1989). *Depth of planting and J-rooting affect Loblolly Pine seedling under stress conditions*. Southern Forest Experiment Station. Dept. Agric. General Technical Report. V74: 169-175.
- BURDETT, A.N.; SIMPSON, D.G.; THOMPSON, C.F. (1983). *Root development and plantation establishment success*. Plant & Soil 71: 103-110.
- BURDETT, A.N. (1987). *Understanding root growth capacity: theoretical considerations in assessing planting stock quality by means of root growth tests*. Can. J. For. Res. V. 17. pp. 768-775.
- BURDETT, A.N. (1990). *Physiological processes in plantation establishment and the development of specifications for forest planting stock*. Can. J. For. Res. 20: 415-427.
- BURNS, R.M. (1974). *Deep planting detrimental to longleaf pines on droughty, acid sands*. Tree Planters' Notes. Fall. 28-29 pp.
- CEBALLOS, L.; RUÍZ DE LA TORRE, J. (1979). *Árboles y arbustos de la España peninsular*. Fundación Conde del Valle de Salazar. ETSIM. Madrid.
- CENTRE TECHNIQUE DU GÉNIE RURAL, DES EAUX ET DES FORÊTS. (1980). *Influence de la position du collet lors de la plantation*. Reveu Forestiere Francaise 32: 6, 561-565.
- CUMMINGS, W. N. (1945). *Cover basin test results for guidance of erosion control planting practice*. Valley Authority. 19 pp. processe.
- DOMÍNGUEZ, S; VILLAR, P; PEÑUELAS, J.L; HERRERO, N; NICOLÁS, J.L; (1999). *Técnicas para cultivar encinas en suelos agrícolas*. Quercus nº 166. pp.22-25
- EGGER B.; EINING W.; SCHLERETH A.; WALLEUDA T.; MAGEL E.; LOEWE A.; HAMPP R. (1996). *Carbohydrate metabolism in one-and two-year-old Spruce needles, and stem carbohydrates from three months before until three months after bud break*. Physiol. Plant. 96: 91-100.
- FERRÁN ARANAZ, M. *SPSS para Windows*. McGraw-Hill. (1996).
- GANDULLO, J. M; et al. (1994). *Estaciones ecológicas de los pinares españoles*. Ecología nº 1. ICONA. Madrid.

GARCÍA SALMERÓN, J.; (1995). *Manual de Repoblaciones Forestales*. Fundación Conde del Valle de Salazar. ETSIM. Madrid.

GEMMEL P., NILSSON U.; WELANDER T. (1996) *Development of oak and beech seedlings planted under varying shelterwood densities and with different site preparation methods in southern Sweden*. New Forests. V 12(2): 141-161.

GIL L.; PRADA M.A. (1993). *Los pinos como especies básicas de la restauración forestal en el medio mediterráneo*. Ecología nº 7: 113-125.

HAYWOOD, J.D.; BARNETT, J.P. (1994). *Comparing methods of artificially regenerating Loblolly and Slash Pines: container planting, bareroot planting and spot seeding*. Tree Planters' Notes. Vol.45(2): 63-67.

HUURI, O. (1972). *The effect of deviating planting techniques on initial development of seedlings of Scots Pine and Norway Spruce*. Communicationes Instituti Forestalis Fenniae 75: 6-92.

HUURI, O. (1978). *Effects of various treatments at planting and of soft containers on the development of scots pine (Pinus sylvestris L.)*. Proceedings of the root Form Planted Trees Symposium, Victoria, British Columbia. Canadian Forestry Service. Join Report nº8: 101-107.

JIMÉNEZ, J.; *et alter*. (19). *Repoblación de montes*. Ministerio de Agricultura. (1956). Madrid.

KAUSHAL, P.; AUSSENAC, G. (1989). *Transplanting shock in Corsican pine and Cedar of Atlas seedlings: Internal water deficits , growth and root regeneration*. Forest Ecology and Management. 27: 29-40.

KOSHI, P. T. (1960). *Deep planting has little effect in a wet year*. U. S. Forest Service. Tree Planters' Notes 40. 24 pp.

KOZLOWSKI, T.; WINGET, C. (1964). *The role of reserves in leaves, branches, stems, and roots on shoot growth of Red pine*. Amer. Jour. Bot. 51(5): 522-529.

LOACH, K.; LITTLE, C.H.A. (1972). *Production, storage, and use of photosynthate during shoot elongation in balsam fir (Abies balsamea)*. Can. J. Bot. 51: 1161-1168.

MATTSSON, A. (1991). *Root growth capacity and field performance of Pinus sylvestris and Picea abies seedlings*. Scand. J. For. Res. 6: 105-112.

McCLAIN, B.; SMITH, Jr. (1976). *Check on longleaf pine depth planting experiment 27 years after planting*. Tree Planters' Notes. 27(2).

McGEE, C.E.; HATCHER, J.B. (1963). *Deep planting Small Slash Pine on old field sites in the Carolina Sandhills*. J. For. 61: 382-383.

NAVARRO, R; PEMÁN GARCÍA, J. *Repoblaciones forestales*. Universitat de Lleida. (1998).

NOLAND, T. L.; MOHAMMED, G. H.; SCOTT, M. (1997). *The dependance of root growth potential on light level, photosynthetic rate, and root starch content in jack pine seedlings*. New Forests 13: 105-119.

PEÑUELAS, J.L; DOMÍNGUEZ, S.; RENILLA, I. (1996). *Experiencias sobre control de la competencia herbácea en repoblaciones de terrenos agrícolas abandonados. Resultados de tres años en campo*. Revista Montes nº 45. pp. 30-36.

PHILIPSON, J. J. (1988). *Root growth in Sitka spruce and Douglas- fir transplants: dependence on the shoot and stored carbohydrates*. Tree Physiology 4: 101-108.

RITCHIE, G.A. (1980). *Root growth potential: its development and espression in forest tree seedlings*. N. Z. J. For. Sci. 10 (1): 218-248.

RITCHIE, G.A. (1982). *Carbohydrate reserves and root growth potential in Douglas-fir seedlings before and after cold storage*. Can. J. For. Res. 12: 905-912.

RITCHIE, G.A. (1985). *Root growth potential: principles, procedures, and predictive ability*. In Proceedings Evaluating seedling quality: principles, procedures, and predictive abilities of major tests. Oregon State University.

ROSE, R. (1992). *Root growth potential and starch differences in seedlings of six families of genetically improved loblolly pine*. Forest Science. Vol. 38. Nº2. pp. 448-456.

ROYO, A.; FERNÁNDEZ, M.; GIL, L.; PARDOS, J.A. (1996). *Predicción de la supervivencia y crecimiento de las plantas de vivero mediante medidas de parámetros fisiológicos pre y post-trasplante*.

SERRADA HIERRO, R.; (1995). *Apuntes de repoblaciones forestales*. Fundación Conde del Valle de Salazar. ETSIM. Madrid.

SHIPMAN, R.D; HATCHER, J.B. (1957). *Planting small slash pine seedlings*. Forest Farmer XVI (5): 7, 18.

SHIVER, B.D.; BORDERS, B.E.; PAGE, H.H.; RAPER, S.M. (1990). *Effect of some seedling morphology and planting quality variables on seedling survival in the Georgia Piedmont*. Southern Journal Applied Forestry 14: 109-115.

SIMPSON, D. G.; RITCHIE, G. A. (1997) *Does RGP predict field performance? A debate*. New Forests 13: 253-277.

SIMPSON, D. G; VYSE, A. (1995). *Planting stock performance: site and RGP effects*. The Forestry Chronicle 71: 739-742.

SIMPSON, D. G.; THOMPSON, C. F.; SUTHERLAND, C. D. (1994). *Field performance potential of interior spruce seedlings: effect of stress treatments and prediction by root growth potential and needle conductance*. Can. J. For. Res. V. 24. pp. 576-586.

SLOCUM, G.K. (1951). *Survival of loblolly pine seedlings as influenced by depth of planting*. Jour. Forestry 49: 500.

SLOCUM, G.K and MAKI, T.E. (1956). *Some effects of depth of planting upon Loblolly Pine in the North Carolina Piedmont*. J. For. Vol. 54. pp. 21-25.

STROTHMAN, R.O. (1976) *Douglas fir seedlings planted by four methods. Results after ten years*. Research Note PSW-310 Pacific Southwest. Forest and Range Experiment Station.

SWITZER, G.L. (1960). *Exposure and planting depth effects on loblolly pine planting stock on poorly drained sites*. J.For. 58: 390-391.

THOMPSON, B.; PUTTONEN, P. (1992). *Patterns of gas exchange, photosynthate allocation, and root growth during a root growth capacity test*. Can. J. For. Res. 22: 248-254.

TINUS, R. W. (1996) *Root growth potential as an indicator of drought stress history*. Tree Physiology 16: 795-799.

VALIO, I.F.M. (1999). *Effects of shading and removal of plant parts on growth of "Trema micrantha" seedlings*. Tree Physiology 21, 65-70.

TSCHAPLINSKI, T. J.; BLAKE, T. J. (1994). *Carbohydrate mobilization following shoot defoliation and decapitation in hybrid poplar*. Tree Physiology 14: 141-451.

VAN DEN DRIESSCHE, R. (1987). *Importance of current photosynthate to new root growth in planted conifer seedlings*. Can. J. For. Res. 17: 776-782.

VAN DEN DRIESSCHE, R. (1991). *New root growth of douglas-fir seedlings at low carbon dioxide concentration*. Tree Physiology 8: 289-295.

VILLAR SALVADOR, P.; OCAÑA, L.; PEÑUELAS, J.; CARRASCO, I.; DOMÍNGUEZ, S. & RENILLA, I. (1997). *Relaciones hídricas y potencial de formación de raíces en plántulas de Pinus halepensis Mill. sometidas a diferentes niveles de endurecimiento por estrés hídrico*. Cuadernos de la S. E. C. F. nº4. pp. 81-92.

WAKELEY, P. C. (1954). *Planting the southern pines*. Southern Forest Experiment Station. U. S. Dept. Agric. Monograph nº 28, 233 pp.